

Topics

- Principals of immunization
- Vaccines
- Immunizations

17-1

Principals of immunization

- Active immunity
- Passive immunity

17-2

	Active	Passive
Natural	Natural exposure to antigen induces an immune response; immunity following an attack of measles.	Transfer of antibodies or cells produced by others; temporary immunity from antibodies of the mother transferred to infant across the placenta or in milk.
Artificial	Deliberate exposure to antigen induces an immune response; immunization of children.	Antibodies in immune serum are introduced into body; injection of rabies immune globulin after a dog bite.

17-3

Figure 17.1 Active and passive immunity

Vaccines

- Attenuated
- Inactivated

17-4

Table 17.1 Some Important Immunizing Agents for Humans

Disease	Type of Vaccine	Persons Who Should Receive the Vaccine
Anthrax	Acellular	People in occupations that put them at risk of exposure, such as military personnel
Diphtheria	Toxoid	Children; adults receive a booster every 10 years
<i>Haemophilus influenzae</i> type b infections	Polysaccharide-protein conjugate	Children
Hepatitis A	Inactivated virus	Children who live in selected regions, people travelling to certain parts of the world
Hepatitis B	Protein subunit produced by genetically engineered <i>Saccharomyces cerevisiae</i> and purified	Children, adults in high-risk groups such as IV drug abusers, health care workers who might be exposed to infected blood, and contacts of infected people, homosexual men, and people who have multiple sexual partners
Influenza	Inactivated virus, usually given by injection in the United States, but as a nasal spray in parts of Europe	Adults over age 50, medical personnel, and people at increased risk for complications; given yearly, as the antigens of the virus change frequently
Measles	Attenuated virus	Children, people entering college, adults born after 1956 who have not been immunized, travelers to foreign countries, and HIV-infected people without severe immunosuppression
Meningococcal disease	Purified polysaccharide (4 serotypes)	Children and adults with certain conditions that put them at greater risk (for example, those without a spleen or who have certain complement system defects); people traveling to sub-Saharan Africa

Table 17.1 - Vaccines

17-5

Table 17.1 Some Important Immunizing Agents for Humans

Disease	Type of Vaccine	Persons Who Should Receive the Vaccine
Mumps	Attenuated virus	Same as measles
Pertussis (whooping cough)	Acellular vaccine given together with diphtheria and tetanus toxoids (DTaP)	Children
Pneumococcal infection	Two forms—purified polysaccharide (PPV) and polysaccharide-protein conjugate (PCV)	Children should receive PCV; adults over 65, people with certain chronic infections, and others in high-risk groups should receive PPV
Rabies	Inactivated virus grown in human or rhesus monkey cells	People exposed to the virus, people at high risk for exposure, such as veterinarians and other animal handlers
Rubella (German measles)	Attenuated virus	Children, adults (particularly women) who are susceptible, health care workers who are at high risk of exposure
Tetanus	Toxoid	Children; adults receive a booster every 10 years
Tuberculosis	Attenuated BCG strain of tuberculosis bacteria	Used only in special circumstances in the United States; widely used in other countries
Typhoid fever	Two forms—attenuated bacteria (taken orally) and purified polysaccharide	People traveling to certain parts of the world
Varicella-zoster (chickenpox)	Attenuated virus	Children; may also be given to susceptible adults
Yellow fever	Attenuated virus	Travelers to affected areas

Table 17.2 - Vaccines

17-6

Immunizations

- Paralytic poliomyelitis
- Effectiveness of immunizations
- Recommended immunizations
- Future immunizations

17-7

Paralytic poliomyelitis

- 1950 – Salk vaccine (inactivated virus)
- 1960 – Sabin vaccine (attenuated virus)
- Salk vaccine is safe
- Sabin vaccine provides herd immunity

17-8

Table 17.3 The Effectiveness of Universal Immunization in the United States

Disease	Cases per Year Before Immunization	Decrease After Immunization
Smallpox	48,164 (1900–1904)	100%
Diphtheria	175,885 (1920–1922)	Nearly 100%
Pertussis (whooping cough)	147,271 (1922–1925)	95.7%
Tetanus	1,314 (1922–1926)	97.4%
Paralytic poliomyelitis	16,316 (1951–1954)	100%
Measles	503,282 (1958–1962)	Nearly 100%
Mumps	152,209 (1968)	99.6%
Rubella (congenital syndrome)	823 (estimated)	99.4%
<i>Haemophilus influenzae</i> type b infections	20,000 (estimated)	99.7%

Table 17.3 - Effectiveness of immunizations

17-9

Table 17.5 Some Diseases for Which New or Improved Vaccines Are Sought

Disease	Estimated impact
HIV/AIDS	40 million infected worldwide, with approximately 14,000 new infections daily
Malaria	300–500 million cases/yr and up to 3 million deaths/yr worldwide
Influenza	30–50 million cases/yr worldwide; 10,000–40,000 deaths/yr in the United States
Strep throat	20 million cases/yr in the United States
Genital herpes	45 million infected and 500,000 new infections/yr in the United States
Hepatitis C	170 million infected worldwide
Cancer	1 in 3 in the United States may get cancer, resulting in 560,000 deaths/yr

Table 17.5 - Future immunizations

17-10


