

Cardiovascular, Pulmonary and Renal Pathology

Homocysteine, System $b^{0,+}$ and the Renal Epithelial Transport and Toxicity of Inorganic Mercury

Christy C. Bridges and Rudolfs K. Zalups

From the Division of Basic Medical Sciences, Mercer University
School of Medicine, Macon, Georgia

Proximal tubular epithelial cells are major sites of homocysteine (Hcy) metabolism and are the primary sites for the accumulation and intoxication of inorganic mercury (Hg^{2+}). Previous *in vivo* data from our laboratory have demonstrated that mercuric conjugates of Hcy are transported into these cells by unknown mechanisms. Recently, we established that the mercuric conjugate of cysteine [2-amino-3-(2-amino-2-carboxy-ethylsulfanylmercurisulfanyl)propionic acid; Cys-S-Hg-S-Cys], is transported by the luminal, amino acid transporter, system $b^{0,+}$. As Cys-S-Hg-S-Cys and the mercuric conjugate of Hcy (2-amino-4-(3-amino-3-carboxy-propylsulfanylmercurisulfanyl)butyric acid; Hcy-S-Hg-S-Hcy) are similar structurally, we hypothesized that Hcy-S-Hg-S-Hcy is a substrate for system $b^{0,+}$. To test this hypothesis, we analyzed the saturation kinetics, time dependence, temperature dependence, and substrate specificity of Hcy-S-Hg-S-Hcy transport in Madin-Darby canine kidney (MDCK) cells stably transfected with system $b^{0,+}$. MDCK cells are good models in which to study this transport because they do not express system $b^{0,+}$. Uptake of Hg^{2+} was twofold greater in the transfectants than in wild-type cells. Moreover, the transfectants were more susceptible to the toxic effects of Hcy-S-Hg-S-Hcy than wild-type cells. Accordingly, our data indicate that Hcy-S-Hg-S-Hcy is transported by system $b^{0,+}$ and that this transporter likely plays a role in the nephropathy induced after exposure to Hg^{2+} . These data are the first to implicate a specific, luminal membrane transporter in the uptake and toxicity of mercuric conjugates of Hcy in any epithelial cell. (*Am J Pathol* 2004, 165:1385–1394)

In recent years, homocysteine (Hcy) has become known as an amino acid of significant clinical importance.^{1,2} Normally, Hcy forms from the intracellular metabolism of methionine and is subsequently broken down by one of two pathways: remethylation or transsulfuration. In the remethylation pathway, a methyl group, from either *N*⁵-

methyltetrahydrofolate or betaine, is transferred to Hcy to reform methionine. Alternatively, Hcy can enter the transsulfuration pathway, where it is broken down into cysteine (Cys) and α -ketobutyrate via the sequential actions of cystathionine- β -synthase and γ -cystathionase.² Alterations in these metabolic pathways may lead to hyperhomocysteinemia, which can contribute to the induction of cardiovascular disease in some humans.³

A number of studies have implicated the kidney as a major site of Hcy metabolism.^{1,2,4–7} Although both pathways involved in the metabolism of Hcy (ie, transsulfuration and remethylation) are used by renal epithelial cells,⁸ the transsulfuration route is the predominant means by which Hcy is degraded in these cells.^{5,6} Moreover, this degradation occurs primarily in the epithelial cells lining the proximal tubule.⁵ Although our understanding of Hcy metabolism within proximal tubular cells is advancing, the mechanisms by which Hcy gains entry in the cytosolic compartments of these cells remain unclear.

Interestingly, proximal tubular cells are also the primary sites where inorganic mercury (Hg^{2+}) accumulates and exerts its toxic effects. Because of the strong electrophilic properties of Hg^{2+} , mercuric ions are carried around in the plasma as conjugates of thiol-containing biomolecules. Extracellular thiols that have been implicated in the binding of Hg^{2+} include Cys and glutathione (GSH). In fact, *in vitro* data from our laboratory indicate that mercuric conjugates of Cys, particularly in the form of 2-amino-3-(2-amino-2-carboxy-ethylsulfanylmercurisulfanyl)propionic acid (Cys-S-Hg-S-Cys), are taken up at the luminal plasma membrane of proximal tubular cells by the heterodimeric amino acid transporter, system $b^{0,+}$.⁹ This transporter is comprised of a light chain subunit, $b^{0,+}$ AT and a heavy chain subunit, rBAT.^{10,11} It should be pointed out that system $b^{0,+}$ is the principal

Supported, in part, by the National Institutes of Health (National Institute of Environmental Health Services grants, ES05980, ES05157, and ES11288, to R.K.Z. and ES012556 to C.C.B.), and the Ruth L. Kirschstein National Research Service Award to C.C.B. ES012556 is the NRSA awarded to C.C.B.

Accepted for publication June 3, 2004.

Address reprint requests to Dr. Rudolfs K. Zalups, Mercer University School of Medicine, Department of Basic Medical Sciences, 1550 College St., Macon, GA 31207. E-mail: zalups.rk@mercer.edu.

sodium-independent amino acid transporter involved in the luminal absorption of the amino acid cystine along the proximal tubule.¹⁰⁻¹⁴ Because of the structural homology between Cys-S-Hg-S-Cys and cystine (Cys-S-S-Cys), we proposed recently that Cys-S-Hg-S-Cys behaves as a molecular mimic of cystine at system b^{0,+}.⁹

Hcy is a thiol-containing amino acid and is a homologue of Cys. Thus one would expect this amino acid to form linear II coordinate covalent complexes with Hg²⁺. Indeed, our laboratory has shown indirectly that the mercuric conjugate of Hcy, 2-amino-4-(3-amino-3-carboxy-propylsulfanylmercuricsulfanyl)butyric acid (Hcy-S-Hg-S-Hcy) is transported by proximal tubular cells.¹⁵ When Hg²⁺ was administered intravenously to rats as Hcy-S-Hg-S-Hcy, some of the administered Hg²⁺ was taken up at the luminal plasma membrane of proximal tubular epithelial cells.¹⁵ The mechanism(s) for this uptake has/have not yet been defined. However, given that Hcy-S-Hg-S-Hcy is similar structurally to the disulfide amino acid homocystine (Hcy-S-S-Hcy), the participation of luminal amino acid transporters seems to be a logical possibility. Some support for this notion comes in part from *in vitro* data showing that the transport of Hcy and homocystine in the kidneys is mediated by a high-affinity transporter with a substrate specificity similar to that characterized for the cystine transporter, system b^{0,+}.¹⁶ Additional evidence from *in vitro* studies using retinal pigment epithelial cells indicates that Hcy is a substrate of system b^{0,+}.¹⁷ As homocystine and cystine are structural homologues, it is logical, therefore, to postulate that homocystine may be transported by the same amino acid transporter (ie, system b^{0,+}) that facilitates the uptake of cystine and the mercuric conjugate, Cys-S-Hg-S-Cys. Moreover, given the structural similarity between Cys-S-Hg-S-Cys and Hcy-S-Hg-S-Hcy, we hypothesize that system b^{0,+} can mediate the uptake of Hcy-S-Hg-S-Hcy in renal epithelial cells expressing this transport system.

We tested this hypothesis by studying and characterizing the transport of Hg²⁺, when presented as Hcy-S-Hg-S-Hcy, in type II Madin-Darby canine kidney (MDCK) cells transfected stably with both subunits of system b^{0,+}. These cells are a line of renal epithelial cells derived from the distal nephron of the dog and were used for the current experiments because they do not normally express b^{0,+}AT or rBAT. Data from the present study indicate clearly that Hg²⁺, in the form of Hcy-S-Hg-S-Hcy, is indeed a transportable, toxic substrate of system b^{0,+}. This study is also the first to provide direct molecular evidence for the participation of a specific amino acid transport system, namely system b^{0,+}, in the absorptive transport of, and cellular intoxication by, Hcy-S-Hg-S-Hcy.

Materials and Methods

Tissue Culture

Stably transfected MDCK cells, strain II, expressing both subunits of system, b^{0,+}, b^{0,+}AT, and rBAT (b^{0,+}AT-rBAT transfectants), were kindly provided by Dr. François Verrey (University of Zurich, Zurich, Switzerland) and were

cultured as described previously.^{9,18} Briefly, cells were cultured in Dulbecco's modified Eagle's medium (Mediatech, Herndon, VA) supplemented with 10% fetal bovine serum (Atlanta Biologicals, Atlanta, GA), 100 U/ml penicillin, 100 µg/ml streptomycin, 1% essential amino acids, 200 µg/ml geneticin, and 150 µg/ml hygromycin B (Invitrogen, Carlsbad, CA). Wild-type cells were cultured in the same media without geneticin or hygromycin B. Cells were passaged by dissociation in 0.25% trypsin (Invitrogen)/0.5 mmol/L ethylenediaminetetraacetic acid in phosphate-buffered saline. Cultures were maintained at 37°C in a humidified atmosphere of 5% CO₂.

Evaluation of Transport

Uptake measurements were performed as described previously with minor changes.^{9,19,20} Both, wild-type and transfected MDCK cells were seeded in 24-well plates at a density of 0.2 × 10⁶ cells/well and were cultured for 24 hours before the experiment. Transfectants were cultured in the presence of 1 µmol/L dexamethasone to induce the expression of rBAT.¹⁸ At the beginning of each experiment, culture media was aspirated and cells were washed with warm uptake buffer (25 mmol/L 4-(2-hydroxyethyl)-1-piperazineethansulfonic acid (HEPES)/Tris, 140 mmol/L *N*-methyl-D-glucamine chloride, 5.4 mmol/L KCl, 1.8 mmol/L CaCl₂, 0.8 mmol/L MgSO₄, and 5 mmol/L glucose, pH 7.5). Uptake was initiated by adding 250 µl of uptake buffer containing radiolabeled substrates. Cells were incubated for 15 minutes at 37°C, unless otherwise stated. Uptake was terminated by aspiration of radiolabeled compounds followed by the addition of ice-cold uptake buffer containing 1 mmol/L sodium 2,3-dimercaptopropane-1-sulfonate (Sigma Chemical Co., St. Louis, MO), a well-known mercury chelator.²¹ Cells were washed twice with 2,3-dimercaptopropane-1-sulfonate and were subsequently solubilized with 1% sodium dodecyl sulfate in 0.2 N NaOH. The cellular lysate was then added to 5 ml of Opti-Fluor scintillation cocktail (Packard Biosciences, Meriden, CT) and the radioactivity contained therein was determined by counting samples in a Beckman LS6500 scintillation counter (Beckman Instruments, Fullerton, CA).

Hcy-S-Hg-S-Hcy was formed by incubating 5 µmol/L HgCl₂, containing ²⁰³Hg²⁺, with 20 µmol/L DL-homocysteine (Sigma Chemical Co.) for 10 minutes at room temperature. The ratio of these compounds ensured that each mercuric ion in solution bonded to the sulfur atom of two molecules of the respective thiol in a linear II coordinate covalent manner. The mercuric conjugates of thiol-containing molecules formed under these conditions have been shown to be thermodynamically stable from a pH of 1 to 14.²²

Transport of Hg²⁺, in the form of Hcy-S-Hg-S-Hcy, was characterized under various conditions. Time-course experiments were performed wherein both transfectants and wild-type cells were incubated with 5 µmol/L Hcy-S-Hg-S-Hcy, containing ²⁰³Hg²⁺, for various periods ranging from 5 to 90 minutes. The saturation kinetics for the transport processes were determined by incubating cells

with Hcy-S-Hg-S-Hcy, containing ²⁰³Hg²⁺, for 15 minutes at 37°C in the presence of unlabeled Hcy-S-Hg-S-Hcy (1, 5, 10, 15, 25, 50, 100 μmol/L). The maximum velocity (V_{max}) and Michaelis-Menten constant (K_m) were calculated using the following formula: $v = (V_{max} \times [S]) / (K_m + [S])$, where v = velocity of transport and $[S]$ = substrate concentration. The temperature dependence for the uptake of Hg²⁺, in the form of Hcy-S-Hg-S-Hcy, was assessed by measuring the uptake of Hg²⁺ at 4°C and 37°C in the presence of unlabeled Hcy-S-Hg-S-Hcy (1, 5, 10, 15, 25, 50 μmol/L). Substrate specificity for the uptake of Hg²⁺ was assessed by incubating cells with Hcy-S-Hg-S-Hcy, containing ²⁰³Hg²⁺, for 15 minutes at 37°C in the presence of amino acids that are substrates of system b⁰⁺ (cystine, Hcy, arginine, leucine, histidine, phenylalanine, lysine, or cycloleucine) or amino acids that are not substrates of this system (glutamate or aspartate). With the exception of cystine, all amino acids were used at a concentration of 3 mmol/L. Because of low solubility, the highest attainable concentration of cystine was 1 mmol/L.

As a negative control, cells were exposed to 5 μmol/L Hg²⁺ in the presence of 20 μmol/L GSH, which promotes the formation of the mercuric conjugate of GSH, G-S-Hg-S-G. Incubation with Cys-S-Hg-S-Cys served as a positive control. These conjugates were formed as described previously for Hcy-S-Hg-S-Hcy. Wild-type and transfected cells were incubated with each conjugate for 30 minutes at 37°C.

To test the ability of Hg²⁺ to bind to competing thiol-containing molecules, cells were incubated with 5 μmol/L Hcy-S-Hg-S-Hcy in the presence of increasing concentrations of GSH (1 to 50 μmol/L). The incubation was performed for 30 minutes at 37°C.

Cystine Efflux Assays

The ability of Hcy-S-Hg-S-Hcy to stimulate the efflux of cystine was measured in wild-type cells and b⁰⁺AT-rBAT transfectants. Cells were exposed to 1 mmol/L cystine, containing [³⁵S]-cystine, for 10 minutes at 37°C. Extracellular cystine was aspirated and the cells were washed with warm uptake buffer. To measure the effects of various compounds on the efflux of [³⁵S]-cystine/Cys (some of the cystine may have been reduced intracellularly), cells were incubated for 1 minute at 37°C with either uptake buffer, 1 mmol/L unlabeled cystine, Cys-S-Hg-S-Cys, Hcy-S-Hg-S-Hcy, or homocystine. After this incubation, the extracellular fluid was removed and placed in vials containing 5 ml of Opti-Fluor scintillation cocktail. The [³⁵S] contained therein was measured using standard methods for liquid scintillation spectrometry. Cells were then washed twice with ice-cold buffer and were solubilized using 1% sodium dodecyl sulfate/0.2 N NaOH. The cellular content of [³⁵S] was determined by liquid scintillation spectrometry.

Assessment of Cellular Viability

The toxicological effects of Hg²⁺, in the form of Hcy-S-Hg-S-Hcy, were quantified using a methylthiazolotetrazo-

lium (MTT) assay as described previously.^{9,23} This assay measures the activity of mitochondrial dehydrogenase via the conversion of MTT (Sigma Chemical Co.) to formazan crystals. Wild-type and transfected MDCK cells were seeded at a density of 0.2×10^6 cells/ml in 96-well culture dishes (200 μl/well). Cells were cultured for 24 hours, washed twice with warm uptake buffer, and then treated with Hcy-S-Hg-S-Hcy (100, 250, 500, 750, or 1000 μmol/L) in uptake buffer containing glutamine. The incubation was performed for 14 hours at 37°C in a humidified atmosphere of 5% CO₂. Subsequently, cells were washed twice with warm uptake buffer and incubated in MTT (0.5 mg/ml) for 2 hours at 37°C in a humidified atmosphere containing 5% CO₂. After this incubation, solubilization buffer (10% Triton X-100, 0.1 N HCl in isopropyl alcohol) was added to each well and the mixture was allowed to incubate for 16 hours at room temperature. Each plate was read at 490 nm in a BioTek μQuant spectrophotometric plate reader (BioTek, Winooski, VT).

Morphologically discernable pathological changes in the transfected and wild-type MDCK II cells exposed to Hcy-S-Hg-S-Hcy were also characterized and semiquantified microscopically. Wild-type and transfected MDCK cells were seeded in chambered coverslips (Nalge Nunc, Naperville, IL) at a density of 0.2×10^6 cells/ml (0.5 ml/chamber). Cells were treated with various concentrations of Hcy-S-Hg-S-Hcy in uptake buffer containing glutamine. This incubation was performed for 14 hours at 37°C. After the exposure to Hcy-S-Hg-S-Hcy, cells were washed with buffer and the culture media was returned to the chamber. Microscopic images were captured immediately using an Olympus IX-70 inverted biological microscope (Olympus, Melville, NY) equipped with Normarsky optics (Nikon, Melville, NY). All observations of cells were performed using ×10 eyepieces and a ×20 planfluor objective. Images were captured with a Nikon DXM 1200 digital camera.

Data Analysis

All transport experiments were repeated at least three times and each measurement was performed in quadruplicate. Data are presented as mean ± SE. Data expressed as a percent were first normalized using the arcsine transformation before applying any parametric statistical analysis. This transformation takes the arcsine of the square root of the decimal fraction of the percent score. Data for each parameter assessed were first analyzed for normality and for homogeneity of variance. After determining that the data for each parameter were distributed normally with the Kolmogorov-Smirnov test and that there was equal variance among the groups of data using the Levene's test, the means for each set of data were evaluated with a two-way analysis of variance. When statistically significant *F*-values were obtained with the analysis of variance, Tukey's multiple comparison posthoc procedure was used to assess differences among the means. A *P* value of <0.05 was considered statistically significant.




Figure 1. Uptake of inorganic mercury (Hg^{2+}) in wild-type and $b^{0,+}$ AT-rBAT-transfected MDCK II cells exposed to $5 \mu\text{mol/L}$ Hg^{2+} , as a mercuric conjugate of Hcy (Hcy-S-Hg-S-Hcy). **A:** Cells were exposed to Hcy-S-Hg-S-Hcy, at 37°C , for time periods ranging from 5 to 90 minutes. Samples were collected at indicated times. **B:** The saturation kinetics of the transport of Hcy-S-Hg-S-Hcy were measured by incubating cells for 15 minutes at 37°C with $5 \mu\text{mol/L}$ Hcy-S-Hg-S-Hcy, containing $^{203}\text{Hg}^{2+}$, in the presence of unlabeled Hcy-S-Hg-S-Hcy (1 to $100 \mu\text{mol/L}$). Results are presented as mean \pm SE. Data represent three experiments performed in quadruplicate. *, Significantly different ($P < 0.05$) from the mean for the corresponding group of wild-type cells. +, Significantly different from the mean for the same cell type at the initial time point or concentration. ++, Significantly different from the mean for the same cell type at each of the preceding time points or concentrations. +++, Significantly different from the mean for the same cell type at each of the preceding time points or concentrations. +++++, Significantly different from the mean for the same cell type at each of the preceding time points.

Results

Transport of Hg^{2+} (as Hcy-S-Hg-S-Hcy)

Time-course analyses show that there was time-dependent uptake of Hg^{2+} in both the wild-type cells and the $b^{0,+}$ AT-rBAT transfectants exposed to $5 \mu\text{mol/L}$ Hcy-S-Hg-S-Hcy (Figure 1A). However, with the exception of the 5-minute time point, the level of uptake of Hg^{2+} in the transfectants was significantly greater than that of the wild-type cells at all times studied.

The saturation kinetics for the uptake of Hg^{2+} during the exposure to Hcy-S-Hg-S-Hcy were analyzed in wild-type and $b^{0,+}$ AT-rBAT-transfected cells (Figure 1B). Concentration-dependent uptake of Hg^{2+} during the expo-




Figure 2. Temperature dependence of the uptake of inorganic mercury (Hg^{2+}), as a conjugate of Hcy (Hcy-S-Hg-S-Hcy), in wild-type and $b^{0,+}$ AT-rBAT-transfected MDCK II cells. Cells were exposed to $5 \mu\text{mol/L}$ Hcy-S-Hg-S-Hcy, containing $^{203}\text{Hg}^{2+}$ in the presence of unlabeled Hcy-S-Hg-S-Hcy (1 to $50 \mu\text{mol/L}$), for 30 minutes at 37°C or 4°C . Results are presented as mean \pm SE. Data represent three experiments performed in quadruplicate. *, Significantly different ($P < 0.05$) from the means for the corresponding groups of wild-type cells at 4°C and 37°C and significantly different from the mean for the corresponding group of transfectants at 4°C . +, Significantly different from the mean for the transfectants at the initial concentration at 37°C . ++, Significantly different from the mean for the transfectants at each of the preceding concentrations at 37°C . +++, Significantly different from the mean for the transfectants at each of the preceding concentrations at 37°C . +++++, Significantly different from the mean for the transfectants at each of the preceding concentrations at 37°C .

sure to Hcy-S-Hg-S-Hcy was significantly greater in the $b^{0,+}$ AT-rBAT transfectants than in the wild-type cells. In the transfectants, the estimated V_{max} for transport was $60.8 \pm 7.0 \text{ pmol} \times \text{mg protein}^{-1} \times \text{minute}^{-1}$, whereas the K_m was calculated to be $144.4 \pm 24.8 \mu\text{mol/L}$.

Temperature-dependent uptake of Hg^{2+} was detected only in the $b^{0,+}$ AT-rBAT transfectants exposed to Hcy-S-Hg-S-Hcy (Figure 2). At 37°C , the uptake of Hg^{2+} , as Hcy-S-Hg-S-Hcy, was significantly greater in the transfectants than in the corresponding groups of wild-type cells. Furthermore, the amount of transport in the transfectants at 37°C was significantly greater than that observed at 4°C . When the experimental temperature was maintained at 4°C , the association of Hg^{2+} with either cell type was minimal. There were no significant differences in the accumulation of Hg^{2+} between corresponding groups of transfectants and wild-type cells. Moreover, the amount of substrate associated with either cell type at 4°C was similar to that associated with wild-type cells at 37°C .

The findings obtained in the substrate specificity experiments for the transport of Hg^{2+} , in the form of Hcy-S-Hg-S-Hcy, are presented in Figure 3. In the transfectants MDCK II cells exposed to Hcy-S-Hg-S-Hcy, the uptake of Hg^{2+} was reduced significantly by the presence of unlabeled Hcy, arginine, leucine, histidine, phenylalanine, lysine, or cycloleucine. The amino acids, glutamate and aspartate, which are not substrates of system $b^{0,+}$, did not significantly alter the transport of Hg^{2+} . Interestingly, co-incubation of Hcy-S-Hg-S-Hcy with unlabeled cystine resulted in a twofold increase in the uptake of Hg^{2+} in the transfectants MDCK cells. This stimulation was also observed at lower concentrations of cystine (Figure 4A).




Figure 3. Substrate specificity analyses of the uptake of inorganic mercury (Hg²⁺) in wild-type and b⁰⁺AT-rBAT-transfected MDCK II cells. Cells were incubated for 15 minutes at 37°C with 5 μmol/L Hg²⁺, as a conjugate of Hcy (Hcy-S-Hg-S-Hcy), in the presence of various unlabeled amino acids (3 mmol/L; unlabeled cystine = 1 mmol/L). Results are presented as mean ± SE. Data represent three experiments performed in quadruplicate. *, Significantly different (*P* < 0.05) from the mean for the corresponding group of wild-type cells. +, Significantly different from the mean for the control group of the corresponding cell type.

Similarly, when the transport of Hg²⁺ was measured in the presence of excess homocystine, uptake of Hg²⁺ was stimulated (Figure 4B). Interestingly, this stimulation occurred in both wild-type and transfected cells.

Alternatively, when the uptake of [³⁵S]-cystine was measured in the presence of excess Hcy-S-Hg-S-Hcy, transport in the b⁰⁺AT-rBAT-transfectants was inhibited (Figure 5A). The IC₅₀ (50% inhibition) for this process was calculated to be 37.2 ± 7.9 μmol/L. There were no significant differences among groups of wild-type cells. The uptake of other substrates for system b⁰⁺, ie, [³H]-arginine or [³H]-lysine, was also inhibited by the presence of excess Hcy-S-Hg-S-Hcy (data not shown). Furthermore, incubation with increasing concentrations of homocystine (Figure 5B) or Hcy (Figure 5C) inhibited the uptake of [³⁵S]-cystine in the transfectants in a dose-dependent manner.

Efflux Assays

Efflux assays were performed in wild-type cells and b⁰⁺AT-rBAT transfectants to determine the ability of unlabeled cystine, Hcy-S-Hg-S-Hcy, Cys-S-Hg-S-Cys, or homocystine to stimulate the efflux of [³⁵S]-cystine (Figure 6A). Cells were preloaded with [³⁵S]-cystine for 10 minutes at 37°C, after which the efflux of [³⁵S] (the radiolabel may be in the form of cystine or Cys as some intracellular reduction may occur) was measured for 1 minute at 37°C in the presence of the aforementioned compounds. There were no significant differences in the efflux of cystine among the treatment groups of wild-type cells. In the transfectants, however, Hcy-S-Hg-S-Hcy, Cys-S-Hg-S-Cys, homocystine, and cystine stimulated the efflux of [³⁵S]. The efflux of [³⁵S] was greatest when cells were treated with Cys-S-Hg-S-Cys (369.7 ± 25 nmol × mg⁻¹ × minute⁻¹) or Hcy-S-Hg-S-Hcy (364.4 ± 20.7 nmol × mg⁻¹ × minute⁻¹). This efflux was signifi-




Figure 4. Uptake of inorganic mercury (Hg²⁺), as a conjugate of Hcy (Hcy-S-Hg-S-Hcy), in the presence of unlabeled cystine (A) (1 to 1000 μmol/L) or unlabeled homocystine (B) (1 to 500 μmol/L) in wild-type and b⁰⁺AT-rBAT-transfected MDCK II cells. Cells were incubated for 30 minutes at 37°C. Results are presented as mean ± SE. Data represent three experiments performed in quadruplicate. *, Significantly different (*P* < 0.05) from the mean for the corresponding group of wild-type cells.

cantly greater than that in corresponding cells exposed to cystine (215.3 ± 8.8 nmol × mg⁻¹ × minute⁻¹) or homocystine (260.9 ± 9.2 nmol × mg⁻¹ × minute⁻¹). The efflux of [³⁵S] was significantly lower when the extracellular fluid consisted only of uptake buffer (119.2 ± 16.2 nmol × mg⁻¹ × minute⁻¹). When the cellular content of [³⁵S] was measured, the pattern of cystine uptake corresponded inversely to the pattern of cystine efflux described above, ie, the greater the efflux, the lower the cellular content of [³⁵S] (Figure 6B).

Transport of Mercuric Conjugates of Thiol-Containing Biological Molecules

The uptake of Hg²⁺, as a mercuric conjugate of Hcy (Hcy-S-Hg-S-Hcy), Cys (Cys-S-Hg-S-Cys), or GSH (G-S-Hg-S-G) was measured in wild-type cells and transfectants (Figure 7A). When the transfected cells were exposed to Hcy-S-Hg-S-Hcy, the uptake of Hg²⁺ was threefold greater than that in the corresponding group of wild-type cells. Incubation with Cys-S-Hg-S-Cys resulted in a twofold increase in the uptake of Hg²⁺ compared




Figure 5. Uptake of cystine in wild-type and $b^{0,+}$ AT-rBAT-transfected MDCK II cells. Cells were exposed to 5 μ mol/L cystine, containing 35 S-cystine, in the presence of unlabeled Hcy-S-Hg-S-Hcy (A), homocysteine (B), or Hcy (C) for 15 minutes at 37°C. Results are presented as mean \pm SE. Data represent three experiments performed in quadruplicate. *, Significantly different ($P < 0.05$) from the mean for the corresponding group of wild-type cells. +, Significantly different from the mean for the same cell type at the initial concentration. ++, Significantly different from the mean for the same cell type at each of the preceding concentrations. +++, Significantly different from the mean for the same cell type at each of the preceding concentrations. +++++, Significantly different from the mean for the same cell type at each of the preceding concentrations.

with that in the corresponding group of wild-type cells. When the cells were exposed to G-S-Hg-S-G, there were no significant differences in the uptake of Hg^{2+} between corresponding groups of wild-type and transfected




Figure 6. Efflux of ^{35}S (A) and cellular content of ^{35}S after efflux (B) in wild-type MDCK II cells and $b^{0,+}$ AT-rBAT-transfectants. Cells were exposed to 1 mmol/L cystine, containing ^{35}S -cystine, for 10 minutes at 37°C and were subsequently incubated with buffer only, 1 mmol/L cystine, homocysteine, or mercuric conjugates of Cys (Cys-S-Hg-S-Cys) or Hcy (Hcy-S-Hg-S-Hcy) for 1 minute at 37°C. Results are presented as mean \pm SE. Data represent three experiments performed in triplicate. *, Significantly different ($P < 0.05$) from the mean for the corresponding group of wild-type cells. +, Significantly different from the mean for the group of transfectants exposed to buffer. ++, Significantly different from the mean for the transfectants exposed to buffer, cystine, or homocysteine. +++, Significantly different from the mean for all other groups of transfected cells.

MDCK II cells. These data confirmed previously reported findings.⁹

Thiol Competition Assays

Thiol competition assays were performed to determine whether GSH was able to compete with Hcy to bind Hg^{2+} . The formation of G-S-Hg-S-G was measured indirectly by incubating wild-type and $b^{0,+}$ AT-rBAT-transfected cells with Hg^{2+} , as Hcy-S-Hg-S-Hcy, in the presence of increasing concentrations of GSH. As the GSH concentration increased, the amount of Hg^{2+} taken up by the transfectants was reduced significantly. There were no significant differences in the accumulation of Hg^{2+} among groups of wild-type cells (Figure 7B).

Cellular Viability Assays

To determine the relationship between cellular transport and intoxication, the cellular viability of wild-type and




Figure 7. Uptake of inorganic mercury (Hg^{2+}) in wild-type and $b^{0,+}$ AT-rBAT-transfected MDCK II cells. **A:** Cells were exposed to $5 \mu\text{mol/L}$ Hg^{2+} as a conjugate of $20 \mu\text{mol/L}$ Hcy (Hcy-S-Hg-S-Hcy), Cys (Cys-S-Hg-S-Cys), or GSH (G-S-Hg-S-G) for 30 minutes at 37°C . **B:** Cells were exposed to $5 \mu\text{mol/L}$ Hcy-S-Hg-S-Hcy, containing $^{203}\text{Hg}^{2+}$, for 30 minutes at 37°C in the presence of increasing concentrations of GSH. Results are presented as mean \pm SE. Data represent three experiments performed in quadruplicate. *, Significantly different ($P < 0.05$) from the mean for the corresponding group of wild-type cells. +, Significantly different from the mean for the corresponding group of transfected cells treated with Hcy-S-Hg-S-Hcy (**A**) or significantly different from the mean of the same cell type at the initial concentration of GSH (**B**).

transfected cells exposed to Hg^{2+} was assessed. After a 14-hour exposure to $50 \mu\text{mol/L}$ Hcy-S-Hg-S-Hcy, the viability of the $b^{0,+}$ AT-rBAT transfectants was reduced by 12%; exposure to 1 mmol/L Hcy-S-Hg-S-Hcy decreased viability by 85% (Figure 8). The cellular viability of the wild-type cells was decreased significantly (20%) only after exposure to 1 mmol/L Hcy-S-Hg-S-Hcy. Treatment with $5 \mu\text{mol/L}$ Hcy-S-Hg-S-Hcy for 30 minutes (experimental conditions used to study various characteristics of transport) did not significantly reduce the cellular viability in either cell type (data not shown).

Morphological Assessment of Toxicity

The appearance of pathological changes was documented in wild-type and transfected cells exposed to various concentrations of Hg^{2+} , as Hcy-S-Hg-S-Hcy. After a 14-hour exposure to $500 \mu\text{mol/L}$ Hg^{2+} , the wild-type cells were confluent and exhibited few discernable mor-




Figure 8. Cellular viability of wild-type MDCK II cells and $b^{0,+}$ AT-rBAT-transfectants after a 14-hour treatment with various concentrations of inorganic mercury (Hg^{2+}), as a conjugate of Hcy (Hcy-S-Hg-S-Hcy). Results are presented as percent control. Data represent two experiments performed in quadruplicate. *, Significantly different ($P < 0.05$) from the mean for the corresponding group of wild-type cells. +, Significantly different from the mean for the same cell type at the initial concentration. **, Significantly different from the mean for the same cell type at each of the preceding concentrations. ***, Significantly different from the mean for the same cell type at each of the preceding concentrations.

phological changes (Figure 9). In contrast, after the Hg^{2+} -exposure, the transfected cells were no longer confluent and many of the cells appeared spherical as they lost their attachments to the culture surface (Figure 9). In addition, many cells contained large vacuoles and exhibited fragmented DNA.

Discussion

Inorganic mercury (Hg^{2+}) binds readily to thiol-containing biomolecules, such as GSH, Cys, and Hcy. These three thiols are present in plasma at similar concentrations (5 to $10 \mu\text{mol/L}$) and thus, may each bind Hg^{2+} . Indeed, GSH and Cys have been implicated in the binding and subsequent transport of Hg^{2+} .²¹ Moreover, Cannon and colleagues^{24,25} have proposed that mercuric conjugates of Cys (Cys-S-Hg-S-Cys), which are similar structurally to the amino acid cystine, are taken up by cystine transporters via a mechanism of molecular mimicry. It is thought that Cys-S-Hg-S-Cys acts as a mimic of cystine at the site of these carriers. In fact, recent data from our laboratory indicate that, indeed, the cystine transporter, system $b^{0,+}$, is involved in the luminal uptake of Cys-S-Hg-S-Cys in proximal tubular cells.⁹

Although Hcy and Cys are in plasma at similar concentrations,²⁶ very little is known about the potential role of Hcy in the renal uptake of Hg^{2+} . Given that Hcy is a structural homologue of Cys, it seems logical to hypothesize that this amino acid is involved in the handling and transport of Hg^{2+} in the kidneys. In fact, previous *in vivo* data from our laboratory provided the first line of experimental evidence implicating Hcy-S-Hg-S-Hcy as a transportable substrate along the renal proximal tubule.¹⁵ The actual mechanism(s) involved in this uptake, however, was/were not defined. Consequently, we tested the hypothesis that Hcy-S-Hg-S-Hcy, in addition to Cys-S-Hg-S-




Figure 9. Morphological analysis of wild-type (A) and $b^{0,+}$ AT-rBAT-transfected (B) MDCK II cells exposed to 500 $\mu\text{mol/L}$ inorganic mercury (Hg^{2+}), as a conjugate of Hcy (Hcy-S-Hg-S-Hcy), for 14 hours at 37°C. All observations of cells were performed using $\times 10$ eyepieces and a $\times 20$ planfluor objective. Images are representative of two separate experiments.

Cys, is a transportable substrate of system $b^{0,+}$. We did this by characterizing the transport and toxicity of Hg^{2+} , in the form of Hcy-S-Hg-S-Hcy, in type II MDCK cells that were or were not stably transfected with both subunits of system $b^{0,+}$, ie, $b^{0,+}$ AT and rBAT. The $b^{0,+}$ AT-rBAT-transfected cells have been characterized previously and have been shown to be a reliable *in vitro* model in which to study system $b^{0,+}$ -mediated transport.^{9,18}

Time-course analyses of the transport of Hcy-S-Hg-S-Hcy in wild-type and $b^{0,+}$ AT-rBAT-transfected cells revealed that the rate of uptake of this conjugate in the transfectants was at least twofold greater than that in the wild-type cells. Analysis of the saturation kinetics for the transport of Hcy-S-Hg-S-Hcy also showed a twofold increase in the magnitude of uptake of Hcy-S-Hg-S-Hcy in the transfectants compared with that in wild-type cells. Because the only apparent difference between the wild-type and transfected MDCK cells was the presence of a functional system $b^{0,+}$ transporter, it can be concluded that this transporter was responsible for the differences in transport between the two cell types studied. As a result, the findings obtained from the transfected cells indicate that Hcy-S-Hg-

S-Hcy is indeed a transportable substrate of system $b^{0,+}$ and that this transporter is capable of mediating the absorptive uptake of this conjugate *in vivo*.

It should be mentioned that a small amount of Hg^{2+} was associated with the wild-type cells when the cells were exposed to Hcy-S-Hg-S-Hcy. It is likely, however, that this association represents nonspecific binding and/or uptake via another mechanism. This conclusion is supported by our data showing temperature-dependent transport in only the transfectants. These data support the hypothesis that the uptake Hcy-S-Hg-S-Hcy in the transfectants is a carrier-mediated process.

Substrate-specificity analyses of transport provide further support for the hypothesis that Hcy-S-Hg-S-Hcy is taken up by system $b^{0,+}$. Substrates that are typically transported by system $b^{0,+}$ inhibited the transport of Hcy-S-Hg-S-Hcy whereas amino acids that are not substrates of this transporter did not affect the uptake of this conjugate. This pattern of inhibition parallels that observed when cystine^{9,18} or Cys-S-Hg-S-Cys,⁹ were used as transportable substrates of system $b^{0,+}$.

Interestingly, when the uptake of Hcy-S-Hg-S-Hcy was assessed in the presence of unlabeled cystine or homocystine, inward transport of Hcy-S-Hg-S-Hcy was stimulated twofold or greater in the transfected cells. This stimulation was observed at all concentrations of cystine or homocystine studied. This phenomenon was specific for the substrate Hcy-S-Hg-S-Hcy because the transport of other substrates of system $b^{0,+}$, such as [³⁵S]-cystine, [³H]-arginine, or [³H]-lysine was inhibited when each of these substrates was co-administered with unlabeled cystine or homocystine. One explanation for the observed stimulatory effect is that the presence of excess cystine or homocystine increases the rate of Hcy-S-Hg-S-Hcy uptake by some presently unknown exchange mechanism.

To test this theory, we studied the efflux of [³⁵S]-cystine/Cys (as some of the cystine may be reduced intracellularly) in wild-type and $b^{0,+}$ AT-rBAT-transfected MDCK cells pre-exposed to 1 mmol/L cystine, containing [³⁵S]-cystine. Subsequent incubation with unlabeled Hcy-S-Hg-S-Hcy, Cys-S-Hg-S-Cys, cystine, or homocystine induced the efflux of [³⁵S] from the $b^{0,+}$ AT-rBAT transfectants. Of the compounds tested, Hcy-S-Hg-S-Hcy and Cys-S-Hg-S-Cys appeared to have the greatest effect on the efflux of [³⁵S]. Cystine and homocystine were also able to stimulate efflux, but to a lesser degree. The efflux of [³⁵S] from cells exposed to uptake buffer was significantly less than that in the cells exposed to other compounds. This efflux may represent the fraction of cystine that diffuses passively out of the cells. Although the implications for these data are unclear, the results indicate that there is a unique interaction between Hcy-S-Hg-S-Hcy and cystine, which promotes the uptake of this conjugate. A similar type of interaction has been shown to occur when Cys-S-Hg-S-Cys was used as a transportable substrate of system $b^{0,+}$.⁹

Pathophysiological measurements, in the form of MTT assays, and detailed microscopic analyses from the present study indicate that the amino acid transporter, system $b^{0,+}$, plays a role in the cellular intoxication in-

duced by exposure to Hcy-S-Hg-S-Hcy. Data from the MTT assays and morphological analyses indicate that cells transfected with system b^{0,+} undergo cellular pathology and death after exposure to lower concentrations of Hcy-S-Hg-S-Hcy than do corresponding wild-type cells. These data and the current transport data indicate that a strong relationship exists between the rates of transport of Hcy-S-Hg-S-Hcy by system b^{0,+} and the induction of cellular injury and death. Therefore, it appears that system b^{0,+} participates in the cellular intoxication induced by Hg²⁺ by promoting the uptake of Hcy-S-Hg-S-Hcy. The intracellular mechanisms involved in the induction of renal cellular injury and death remain undefined.

Thiol competition experiments were performed to test indirectly the phenomenon of thiol exchange. The electrophilic properties of Hg²⁺ are such that this metal binds readily to thiol-containing molecules.²¹ The idea of thiol competition would imply that mercuric ions bind preferentially to the most prevalent thiol.²⁷ Indeed, the current data suggest that as the concentration of excess thiol (in this case, GSH) increases, the formation of mercuric conjugates of that thiol (G-S-Hg-S-G) is favored even when Hg²⁺ exists initially as a conjugate of Hcy. This phenomenon of thiol competition would be predicted to occur in cases of hyperhomocysteinemia, in which elevated concentrations of plasma Hcy (as high as 200 μmol/L)^{28,29} would promote the preferential formation of Hcy-S-Hg-S-Hcy after exposure to Hg²⁺. Because this conjugate is a highly transportable species of Hg²⁺, hyperhomocysteinemia may increase the risk of renal injury after exposure to this toxic metal. Interestingly, *in vivo* studies in rats have shown that acute hyperhomocysteinemia results in a fourfold increase in the renal uptake of Hcy.⁶ Therefore, hyperhomocysteinemia not only favors the formation of Hcy-S-Hg-S-Hcy, but also may potentially up-regulate the uptake of Hcy. Given that this conjugate and the amino acid Hcy are both transportable substrates of system b^{0,+}, we can speculate that the probable up-regulation of this system, as may be the case under hyperhomocysteinemic conditions, would not only promote the uptake of Hcy, but also that of a toxic species of Hg²⁺, Hcy-S-Hg-S-Hcy.

In conclusion, the results from the current study demonstrate for the first time that Hg²⁺, in the form of Hcy-S-Hg-S-Hcy, is a transportable substrate of system b^{0,+}. Accordingly, this system may play a role in the nephropathy induced after exposure to Hg²⁺. These results also implicate a mechanism of molecular mimicry whereby Hcy-S-Hg-S-Hcy mimics the amino acid cystine or homocystine at the site of system b^{0,+} to gain access to the intracellular compartment of renal epithelial cells. The data provide strong evidence for the involvement of the Na⁺-independent transporter, system b^{0,+}, in the absorptive transport of Hcy-S-Hg-S-Hcy at the luminal plasma membrane of proximal tubular cells. However, this transporter is likely not the sole mechanism responsible for this transport. Previous studies in isolated perfused renal tubules have provided strong evidence for the involvement of at least one Na⁺-dependent transport system.²⁵ Additional studies are clearly needed to provide a more complete understanding of the mechanisms

by which Hg²⁺ is transported across the luminal plasma membranes of proximal tubular epithelial cells.

Acknowledgments

We thank Dr. Delon Barfuss at Georgia State University, Atlanta, GA, for his assistance with the manufacture of the ²⁰³Hg; Dr. François Verrey at the University of Zurich, Zurich, Switzerland, for kindly providing the b^{0,+}AT-rBAT-transfected MDCK II cells; and Ms. Jamie Battle for technical assistance.

References

1. Refsum H, Ueland PM, Nygard O, Vollset SE: Homocysteine and cardiovascular disease. *Annu Rev Med* 1998, 49:31–62
2. Selhub J: Homocysteine metabolism. *Annu Rev Nutr* 1999, 19:217–246
3. Clarke R, Daly L, Robinson K, Naughten E, Cahalane S, Fowler B, Graham I: Hyperhomocysteinemia: an independent risk factor for vascular disease. *N Engl J Med* 1991, 324:1149–1155
4. Dudman NP, Guo XW, Gordon RB, Dawson PA, Wilcken DE: Human homocysteine catabolism: three major pathways and their relevance to development of arterial occlusive disease. *J Nutr* 1996, 126:295S–300S
5. House JD, Brosnan ME, Brosnan JT: Characterization of homocysteine metabolism in the rat kidney. *Biochem J* 1997, 328:287–292
6. House JD, Brosnan ME, Brosnan JT: Renal uptake and excretion of homocysteine in rats with acute hyperhomocysteinemia. *Kidney Int* 1998, 54:601–607
7. Bostom A, Brosnan JT, Hall B, Nadeau MR, Selhub J: Net uptake of plasma homocysteine by the rat kidney *in vivo*. *Atherosclerosis* 1995, 116:59–62
8. Friedman AN, Bostom AG, Selhub J, Levey AS, Rosenberg IH: The kidney and homocysteine metabolism. *J Am Soc Nephrol* 2001, 12: 2181–2189
9. Bridges CC, Bauch C, Verrey F, Zalups RK: Mercuric conjugates of cysteine are transported by the amino acid transporter, system b^{0,+}: implications of molecular mimicry. *J Am Soc Nephrol* 2004, 15:663–673
10. Palacin M, Estevez R, Bertran J, Zorzano A: Molecular biology of plasma membrane amino acid transporters. *Physiol Rev* 1998, 78: 969–1054
11. Palacin M, Fernandez E, Chillaron J, Zorzano A: The amino acid transporter system b^{0,+} and cystinuria. *Mol Membr Biol* 2001, 18: 21–26
12. Kanai Y, Stelzner MG, Lee WS, Wells RG, Brown D, Hediger MA: Expression of mRNA (D2) encoding a protein involved in amino acid transport in S3 proximal tubule. *Am J. Physiol* 1992, 263:F1087–F1092
13. Furriols M, Chillaron J, Mora C, Castello A, Bertran J, Camps M, Testar X, Vilaro S, Zorzano A, Palacin M: rBat, related to L-cystine transport, is localized to the microvilli of proximal straight tubules, and its expression is regulated in kidney by development. *J Biol Chem* 1993, 268:27060–27068
14. Pfeiffer R, Loffing J, Rossier G, Bauch C, Meier C, Eggemann T, Loffing-Cueni D, Kuhn LC, Verrey F: Luminal heterodimeric amino acid transporter defective in cystinuria. *Mol Biol Cell* 1999, 10:4135–4147
15. Zalups RK, Barfuss DW: Participation of mercuric conjugates of cysteine, homocysteine, and N-acetylcysteine in mechanisms involved in the renal tubular uptake of inorganic mercury. *J Am Soc Nephrol* 1998, 9:551–561
16. Foreman JW, Wald H, Blumberg G, Pepe LM, Segal S: Homocysteine uptake in isolated rat renal cortical tubules. *Metabolism* 1982, 31: 613–619
17. Naggar H, Fei YJ, Ganapathy V, Smith SB: Regulation of reduced-folate transporter-1 (RFT-1) by homocysteine and identity of transport

- systems for homocysteine uptake in retinal pigment epithelial (RPE) cells. *Exp Eye Res* 2003, 77:687–697
18. Bauch C, Verrey F: Apical heterodimeric cystine and cationic amino acid transporter expressed in MDCK cells. *Am J Physiol* 2003, 283: F181–F189
 19. Bridges CC, Kekuda R, Wang H, Prasad PD, Mehta P, Huang W, Smith SB, Ganapathy V: Structure, function, and regulation of human cystine/glutamate transporter in retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci* 2000, 42:47–54
 20. Bridges CC, Ola S, Prasad PD, El-Sherbeny A, Ganapathy V, Smith SB: Regulation of taurine transporter expression by NO in cultured human retinal pigment epithelial cells. *Am J Physiol* 2000, 281:C1825–C1836
 21. Zalups RK: Molecular interactions with mercury in the kidney. *Pharmacol Rev* 2000, 52:113–143
 22. Fuhr B, Rabenstein DL: Nuclear magnetic-resonance studies of solution chemistry of metal-complexes. 9. Binding of cadmium, zinc, lead, and mercury by glutathione. *J Am Chem Soc* 1973, 95:6944–6950
 23. Aslamkhan AG, Han YH, Yang XP, Zalups RK, Pritchard JB: Human renal organic anion transporter 1-dependent uptake and toxicity of mercuric-thiol conjugates in Madin-Darby canine kidney cells. *Mol Pharmacol* 2003, 63:590–596
 24. Cannon VT, Barfuss DW, Zalups RK: Molecular homology and the luminal transport of Hg^{2+} in the renal proximal tubule. *J Am Soc Nephrol* 2000, 11:394–402
 25. Cannon VT, Zalups RK, Barfuss DW: Amino acid transporters involved in luminal transport of mercuric conjugates of cysteine in rabbit proximal tubule. *J Pharmacol Exp Ther* 2001, 298:780–789
 26. Ueland PM, Refsum H, Stabler SP, Malinow MR, Andersson A, Allen RH: Total homocysteine in plasma or serum. Methods and clinical applications. *Clin Chem* 1993, 39:1764–1779
 27. Rabenstein DL, Fairhurst MT: Nuclear magnetic resonance studies of the solution chemistry of metal complexes XI. The binding of methylmercury by sulfhydryl-containing amino acids and by glutathione. *J Am Chem Soc* 1975, 97:2086–2092
 28. Stabler SP, Marcell PD, Podell ER, Allen RH, Savage DG, Lindenbaum J: Elevation of total homocysteine in the serum of patients with cobalamin or folate deficiency detected by capillary gas chromatography-mass spectrometry. *J Clin Invest* 1988, 81:466–474
 29. Malinow MR: Hyperhomocysteinemia: a common and easily reversible risk factor of occlusive atherosclerosis. *Circulation* 1990, 81: 2004–2006